



Record of immunisation QI activity

| PIP QI quarter: <i>please tick</i> ✓ | Quarter 1 | Quarter 2 | Quarter 3 | Quarter 4 |
|--|--|---|---|---|
| Date: | <i>Nov – Jan</i> <input type="checkbox"/> | <i>Feb – Apr</i> <input type="checkbox"/> | <i>May – Jul</i> <input type="checkbox"/> | <i>Aug – Oct</i> <input type="checkbox"/> |
| QI Activity: | Respiratory syncytial virus (RSV) vaccination uptake in pregnant women. | | | |
| Activity goal: <i>What to improve and timeframe</i> | Increase the proportion of active pregnant women receiving RSV vaccination between 28 and 36 weeks of pregnancy within the next 3 months. | | | |
| Activity measures: <i>What data is used to monitor progress</i> | Use data extraction tool* to measure the percentage of active pregnant women who haven't received their maternal RSV vaccination prior to activity. Extract data monthly to monitor progress. | | | |
| Initial benchmark: <i>Baseline data prior to QI activity</i> | Our data shows we currently have % pregnant women who have not had their maternal RSV vaccination. | | | |
| Activity overview: <i>Action plan to drive improvement</i> | <ol style="list-style-type: none"> 1. Use data extraction tool to find the percentage of active pregnant women who have not had their maternal RSV vaccination. 2. Create patient list of pregnant women who have not had their maternal RSV vaccination. 3. Send patient reminders in accordance with practice policy. 4. Display posters on pertussis vaccination in pregnancy in waiting room and have brochures available. 5. Discuss QI activity at staff meetings. 6. Review data monthly to track progress. | | | |
| 30-day checkpoint: <i>Monitor progress at 30 days</i> | | | | |
| 60-day checkpoint: <i>Monitor progress at 60 days</i> | | | | |
| Final 90-day checkpoint: <i>Results at 90 days – summarise improvement from baseline</i> | | | | |
| Reflection: <i>Reflect on achievements, challenges and lessons</i> | | | | |

*Data extraction tools include: PenCAT, POLAR, clinical software reports, etc