







Record of immunisation QI activity

PIP QI quarter: please tick ✓	Quarter 1 Nov – Jan □	Quarter 2 Feb – Apr □	Quarter 3 May – Jul □	Quarter 4 Aug – Oct □
QI Activity:	Pertussis (whooping cough) vaccination uptake in pregnant women.			
Activity goal: What to improve and timeframe	Increase the proportion of active pregnant women receiving pertussis vaccination between 20 and 32 weeks of pregnancy within the next 3 months.			
Activity measures: What data is used to monitor progress	Use data extraction tool* to measure the percentage of active pregnant women who haven't received their maternal pertussis vaccination prior to activity. Extract data monthly to monitor progress.			
Initial benchmark: Baseline data prior to QI activity	Our data shows we currently have % pregnant women who have not had their maternal pertussis vaccination.			
Activity overview: Action plan to drive improvement	 Use data extraction tool to find the percentage of active pregnant women who have not had their maternal pertussis vaccination. Create patient list of pregnant women who have not had their maternal pertussis vaccination. Send patient reminders in accordance with practice policy. Display posters on pertussis vaccination in pregnancy in waiting room and have brochures available. Discuss QI activity at staff meetings. Review data monthly to track progress. 			
30-day checkpoint: Monitor progress at 30 days				
60-day checkpoint: Monitor progress at 60 days				
Final 90-day checkpoint: Results at 90 days – summarise improvement from baseline				
Reflection: Reflect on achievements, challenges and lessons				

*Data extraction tools include: PenCAT, POLAR, clinical software reports, etc

