



Record of immunisation QI activity

PIP QI quarter: <i>please tick</i> ✓	Quarter 1	Quarter 2	Quarter 3	Quarter 4
Date:	<i>Nov – Jan</i> <input type="checkbox"/>	<i>Feb – Apr</i> <input type="checkbox"/>	<i>May – Jul</i> <input type="checkbox"/>	<i>Aug – Oct</i> <input type="checkbox"/>
QI Activity: AIR010A report	Overdue pneumococcal vaccination in adults aged 70–80 years.			
Activity goal: <i>What to improve and timeframe</i>	To decrease the number of patients aged 70–80 years who are overdue for pneumococcal vaccination from baseline within 3 months.			
Activity measures: <i>What data is used to monitor progress</i>	Generate an AIR report listing patients aged 70–80 years who are overdue for pneumococcal vaccination before QI activity commencement, and rerun report when the activity is complete to measure improvement.			
Initial benchmark: <i>Baseline data prior to QI activity</i>	AIR report shows that there are _____ number of patients aged 70-80 years who are overdue for pneumococcal vaccination.			
Activity overview: <i>Action plan to drive improvement</i>	<ol style="list-style-type: none"> 1. Generate a list of patients aged 70–80 years who are overdue for their pneumococcal vaccination via AIR010A report from the Australian Immunisation Register. Use report parameters specific for this cohort. 2. Contact patients to offer free catch-up pneumococcal vaccinations, funded on the NIP for people aged 70 years and over. 3. Display pneumococcal vaccination posters in waiting room and have brochures available. 4. Discuss at staff meetings. 5. Review data monthly to track progress. 			
30-day checkpoint: <i>Monitor progress at 30 days</i>				
60-day checkpoint: <i>Monitor progress at 60 days</i>				
Final 90-day checkpoint: <i>Results at 90 days – summarise improvement from baseline</i>				
Reflection: <i>Reflect on achievements, challenges and lessons</i>				

*Data extraction tools include: PenCAT, POLAR, clinical software reports, etc