When vaccinating young children – Remember to check: Spikevax (Moderna) 6 months – 5 years

Vaccine Preparation		Vaccine Administ	
Rememb	er to check:	Remembe	er to ch
Thaw expiry date	30 days from thaw date	Check the Australian Immur child's full immunisation histor	
Manufacture expiry date	Check current guidelines	Consent form completed?	Mode
Time vial was first punctured	Up to 6 hours (up to 25°C)	Right vaccine for the right patient?	• 6 m
Time the syringe was drawn up	Within 6 hours (if stored between 2°C to 8°C) OR within 1 hour (if stored between 8°C to 30°C) Drawn up doses should not be stored above 30°C	Inght patient:	 Dat Are imr Do Cor cor risk
Vaccine type	Blue/puprle - pediatric formula	Right dose for this patient and cohort?	0.25m (25µg
DISCARD the vaccine if it of	exceeds expiry dates and times	Right dose interval?	8 wee
DISCARD the vaccine if it exceeds expiry dates and times. Checking AIR before administering the vaccine will help avoid vaccination administration errors.		Has the vial been diluted (if required)?	DO NO

*Previous vaccination doses and their brand; the Australian Technical Advisory Group on Immunisation (ATAGI) do not recommend the use of mixed primary schedules in this age group Booster doses are not currently recommended for patients in this age group <u>ATAGI Recommendations on the use of a booster dose of COVID-19 vaccine</u>

> If you respond 'No' to any of the above, DO NOT administer the vaccine. Contact the Vaccine Operations Centre for further advice.

stration

check: 📀

n Register (AIR) for the ore administering vaccine*

lerna 6 months - 5 years sent form

months - 5 years of age ate of birth checked

re they severely

nmunocompromised

o they have a disability

Complex and/or multiple health

onditions which increase the

sk of severe COVID-19

iml 1g per dose)

eeks

NOT DILUTE

